

## King's Research Portal

DOI:

[10.1016/j.jid.2018.05.013](https://doi.org/10.1016/j.jid.2018.05.013)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Pigors, M., Common, J. E. A., Wong, X. F. C. C., Malik, S., Scott, C. A., Tabarra, N., Liany, H., Liu, J., Limviphuvadh, V., Maurer-Stroh, S., Tang, M. BY., Lench, N., Margolis, D. J., van Heel, D. A., Mein, C. A., Novak, N., Baurecht, H., Weidinger, S., McLean, W. H. I., ... Kelsell, D. P. (2018). Exome Sequencing and Rare Variant Analysis Reveals Multiple Filaggrin Mutations in Bangladeshi Families with Atopic Eczema and Additional Risk Genes. *Journal of Investigative Dermatology*, 138(12), 2674-2677.  
<https://doi.org/10.1016/j.jid.2018.05.013>

### Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.


### General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### Take down policy

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



**Figure 1: Filaggrin (*FLG*) risk variants identified in the Bangladeshi atopic eczema (AE) cohort and novel risk genes.**

**(a)** Schematic representation of *FLG* showing all loss-of-function variants identified in this study. (Red arrows indicate previously unreported variants, variants shown black and gray were identified using whole exome sequencing or targeted re-sequencing, respectively) **(b)**

Exome and targeted resequencing demonstrated an intrafamilial heterogeneity of *FLG* risk variants in the Bangladeshi AE cohort, segregating, in part, within families. Of note, the parents and the oldest sibling (all unaffected) of both families were born in Bangladesh. All other siblings were born in the UK. (Black symbols indicate affected individuals, gray circle indicates a previously affected daughter, white symbols represent unaffected individuals) (c) Heat Maps showing the number of rare variants identified in the different AE cohorts from this study. Data are based on allele frequencies from ethnically-matching subpopulations obtained from the genomic database ExAC. Recessive and dominant variants were considered rare if they had an allele frequency  $<0.05$  and  $<0.01$ , respectively. Filaggrin (*FLG*) loss-of-function variants were excluded in the African-American cohort as described previously (Margolis et al., 2014).